CONTENTS

7TH INTERNATIONAL SYMPOSIUM ON HIGH-PERFORMANCE LIQUID CHROMATOGRAPHY OF PROTEINS, PEPTIDES AND POLYNUCLEOTIDES, WASHINGTON, DC, NOVEMBER 2–4, 1987, PART II

Immunological-chromatographic analysis
by L. J. Janis and F. E. Regnier (West Lafayette, IN, U.S.A.) 1

Isolation of specific lymphocyte receptors by high-performance immunoaffinity chromatography
by T. M. Phillips and S. C. Frantz (Washington, DC, U.S.A.) 13

Use of avidin-coated glass beads as a support for high-performance immunoaffinity chromatography
by J. V. Babashak (Vineland, NJ, U.S.A.) and T. M. Phillips (Washington, DC, U.S.A.) 21

High-performance concanavalin A affinity chromatography of liver and hepatoma membrane proteins
by D. Josić, W. Hofmann, R. Habermann and W. Reutter (Berlin, F.R.G.) 29

Immunoaffinity isolation of the Lewis blood group antigens
by S. C. Frantz, H. F. Henriques and T. M. Phillips (Washington, DC, U.S.A.) . . . 41

High-resolution chromatography of nucleic acids on the Gen-Pak FAX column
by D. J. Stowers, J. M. B. Keim, P. S. Paul and Y. S. Lyoo (Ames, IA, U.S.A.), M. Merion (Milford, MA, U.S.A.) and R. M. Benbow (Ames, IA, U.S.A.) 47

Preparation and chromatographic use of 5′-fluorescent-labelled DNA probes
G. Tous and J. Fausnaugh (Piscataway, NJ. U.S.A.), P. Vieira (Metuchen, NJ, U.S.A.) and S. Stein (Piscataway, NJ, U.S.A.) 67

Purification of transfer RNA species by single-step ion-exchange high-performance liquid chromatography
by R. H. Guenther, D. H. Gopal and P. F. Agris (Columbia, MO, U.S.A.) 79

Synthesis and purification of deoxyribose analogues of NAD^+ by affinity chromatography and strong-anion-exchange high-performance liquid chromatography
by R. Alvarez-Gonzalez (Ardmore, OK, U.S.A.) 89

Staphylococcal nuclease high-performance liquid chromatographic characterization of diaminooctane-modified DNA and its biotin and fluorescein derivatives
D. J. Cecchini, K. L. Guan and R. W. Giese (Boston, MA, U.S.A.) 97

Separation of human factor X from factor Xa by reversed-phase high-performance liquid chromatography
by K. T. Shitanishi and S. W. Herring (Los Angeles, CA, U.S.A.) 107

Application of Marfey's reagent in racemization studies of amino acids and peptides
by G. Szókán, G. Mezö and F. Hudecz (Budapest, Hungary) 115

High-performance liquid chromatographic analysis of arginine-containing peptides in biological fluids by means of a selective post-column reaction with fluorescence detection
by G. R. Rhodes and V. K. Boppana (King of Prussia, PA, U.S.A.) 123

Purification of nucleotide-linked peptide
by V. N. Pandey and M. J. Modak (Newark, NJ, U.S.A.) 133

Determination of the molecular size distribution of immunoglobin G (IgG) in intravenous IgG–albumin formulations by high-performance liquid chromatography
by J. K. Lee, F. J. Deluccia and E. L. Kelly (Horsham, PA, U.S.A.), C. Davidson (Pearl River, NY, U.S.A.) and F. R. Borger (Kankakee, IL, U.S.A.) 141

Comparison of Mono Q, Superose-6, and ABx fast protein liquid chromatography for the purification of IgM monoclonal antibodies
by F.-M. Chen, G. S. Naeve and A. L. Epstein (Los Angeles, CA, U.S.A.) 153

High-performance liquid chromatography and photodiode-array detection of the human protein HC (human complex-forming glycoprotein heterogeneous in charge), a chromophore-associated protein
by J. Escribano (Madrid, Spain), R. Matas (Barcelona, Spain) and E. Méndez (Madrid, Spain) 165

Reversed-phase high-performance liquid chromatography of fertirelin acetate and related compounds
by P. A. Hartman and J. D. Stodola (Kalamazoo, MI, U.S.A.) 177

Nucleotide extraction and quantitation from tomato roots by high-performance liquid chromatography
by I. Dutta, G.A. O'Donovan and D. W. Smith (Denton, TX, U.S.A.) 183

Reversed-phase liquid chromatography and sodium dodecyl sulfate polyacrylamide gel electrophoresis characteristics of a recombinant DNA derived malaria antigen
by K. Benedek, B, Hughes, M. B. Seaman and J. K. Swadesh (King of Prussia, PA, U.S.A.) 191

Semi-preparative purification of recombinant human renin and prorenin
by C. T. Carilli, L. Cameron Wallace, L. M. Smith, M. A. Wong and J. A. Lewicki (Mountain View, CA, U.S.A.) 203

Analysis of allergenic components of a *Parietaria judaica* pollen extract by chromatographic methods for the evaluation of purification procedures
by A. Bassoli, F. Chioccara, G. Di Gregorio, B. Rindone, S. Tollari, P. Flagiani, G. Riva and E. Bolzacchini (Milan, Italy) 209

Purification of cytochrome *c* oxidase from corn root mitochondria by hydrophobic interaction chromatography
by N. Parris, H. A. Gruber, S. M. Mozersky and R. A. Barford (Philadelphia, PA, U.S.A.) 219

High-performance hydrophobic interaction chromatography as a means of identifying estrogen receptors expressing different binding domains
by S. M. Hyder and J. L. Wittliff (Louisville, KY, U.S.A.) 225

Optimal resolution of eye lens γ-crystallins by cation-exchange high-performance liquid chromatography on SynChropak CM300
by R. J. Siezen and E. D. Kaplan (Cambridge, MA, U.S.A.) 239

Separation and characterization of two forms of acetohydroxy acid synthase from Black Mexican Sweet corn cells
by B. K. Singh, M. A. Stidham and D. L. Shaner (Princeton, NJ, U.S.A.) 251

Scale-up from analytical to preparative reversed-phase high-performance liquid chromatography of soybean glycinin
by R. E. Peterson and W. J. Wolf (Peoria, IL, U.S.A.) 263

Hydrophobic interaction chromatography of peptides as an alternative to reversed-phase chromatography
by A. J. Alpert (Columbia and Baltimore, MD, U.S.A.) 269

Optimization of concentration overload in preparative liquid chromatography
by S. Ghodbane and G. Guiochon (Knoxville and Oak Ridge, TN, U.S.A.) 275

Large-scale chromatography of recombinant proteins
by E. Hochuli (Basle, Switzerland) 293

Preparative separation of peptide and protein samples by high-performance liquid chromatography with gradient elution. I. The Craig model as a basis for computer simulations
by L. R. Snyder (Orinda, CA, U.S.A.) and G. B. Cox and P. E. Antle (Wilmington, DE, U.S.A.) 303

Preparative separation of peptide and protein samples by high-performance liquid chromatography with gradient elution. II. Experimental examples compared with theory
by G. B. Cox and P. E. Antle (Wilmington, DE, U.S.A.) and L. R. Snyder (Orinda, CA, U.S.A.) 325

Reversed-phase preparative chromatography of [D-Ala^1]peptide T amide
by M. Knight (Washington, DC, U.S.A.), G. D. Mack and R. Perkins (Rockville, MD, U.S.A.) and T. R. Burke, Jr. (Washington, DC, U.S.A.) 345

Preparative purification of peptides by reversed-phase chromatography. Sample displacement mode *versus* gradient elution mode
by R. S. Hodges, T. W. L. Burke and C. T. Mant (Alberta, Canada) 349

Author Index 363

In articles with more than one author, the name of the author to whom correspondence should be addressed is indicated in the article heading by a 6-pointed asterisk (*)